
MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)

MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)

MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)

MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)

MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)

MicroCLAVE[®] Neutral Connector

Use aseptic technique. Single use only - Do not resterilize.

DIRECTIONS:

1. Prime the MicroCLAVE[®] by inverting to expel air and attach to patient's catheter.
2. To access: swab MicroCLAVE[®], then push and twist administration device into MicroCLAVE[®] until secure.
3. Flush MicroCLAVE[®] after each use with normal saline or in accordance with facility protocol.

Latex Free, DEHP Free

CAUTIONS:

- Federal (USA) law restricts this device to sale by or on the order of a physician.
- MicroCLAVE[®] should be changed in accordance with current, recognized guidelines of IV therapy.
- The MicroCLAVE[®] is compatible with luers having an internal diameter (ID) greater than 0.061" (1.55 mm).

MicroCLAVE[®] is a registered trademark of ICU Medical, Inc. K-45703-118A (1/08)