

Pressure Injectable Arrowg+ard Blue Plus[®] Two-Lumen CVC

Contents:

- 1: Arrowg+ard Blue Plus®1 Two-Lumen Catheter: 7 Fr. (2.5 mm OD) x 20 cm, Pressure Injectable
- Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer 1: Needle
- Injection Needle: SafetyGlide^{™2} 25 Ga. x 1" (2.54 cm) 1.
- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) 1:
- Pressure Transduction Probe 1:
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: SafetyGlide^{™2} 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- Syringe: 3 mL Luer-Lock 1.
- 1: Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm
- 5 mL 1% Lidocaine HCI Solution and Alcohol Prep 1:
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with 1: Hi-Lite Orange™ Tint
- SecondSite[™] Adjustable Hub: Fastener
- SecondSite™ Adjustable Hub: Catheter Clamp 1.
- SharpsAway® Disposal Cup 1:
- SharpsAway® II Locking Disposal Cup 1:
- Drape: 41" x 55" with 4" fenestration 1.
- Needle Holder
- Safety Scalpel: #11 1.
- Checklist/CLIP Sheet 1:
- 1: Sterile Procedure Sign

±Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants. Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F)

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.



- 1: Flow Rate Information Card with Injection Log
- 1: Medication Label: 1% Lidocaine
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm)
- 1: Dressing: Tegaderm^{®4} 10 cm x 12 cm
- Dressing: BIOPATCH®5 1:
- Suture: 3-0 Silk with Curved Needle 1: 1: Tubing±: 20" (50 cm) Extension
- 1: HemoHopper® Fluid Receptacle
- ¹Licensed under US Patent No. 7.329.412.
- ²A trademark of Becton, Dickinson and Company
- 3A registered trademark of CareFusion or one of its subsidiaries.
- ⁴A registered trademark of 3M Company.
- 5A registered trademark of Johnson & Johnson Corporation.

6 Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (14 Ga.)	0.57	4731	10
Proximal (18 Ga.)	0.40	913	10

* Priming volumes are approximate and are done without accessories.

+ Flow rate values are approximate and are determined using deionized water at 100 cm head height.

** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.

LBL052789 R00 (2021-11)