



PEF PK-01552-WS

5^{Fr.} 2^{Lumen}

55 cm catheter length 1018 inch dia. spring-wire guide

Two-Lumen PICC

Contents:

- 1: Two Lumen Catheter: 5 Fr. (1.76 mm OD) x 55 cm, T-Port Connector, Blue FlexTip®, Contamination Guard and Placement Wire
- Two Lurieri Garieter. 3 Tr. (1.7.4 mills 2.7.1 mills

- Dilator

 Spring-Wire Guide: .018" (0.46 mm) dia. x 13-1/8" (33 cm) (Straight Soft Tip on Both Ends)

 Catheter: 22 Ga. x 1-3/4" (4.45 cm) Radiopaque over 25 Ga. RW Introducer Needle
 Injection Needle: 18 Ga. x 1-1/2" (3.81 cm)
 Introducer Needle: 21 Ga. x 1-1/2" (3.81 cm)
 Introducer Needle: Echogenic 21 Ga. x 2-3/4" (7 cm) TW
 Introducer Needle: Echogenic 21 Ga. x 2-3/4" (7 cm) TW
 Syringe: 10 mL Luer-Slip
 Skin Protectant Prep
 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®¹ One-Step Solution with Hi-Lite Orange™ Tint
 Dust Cap: Non-Vented
 SecondSite™ Adjustable Hub: Fastener
 SecondSite™ Adjustable Hub: Catheter Clamp
 SharpsAway® Disposal Cup
 Towel

- Towel: 24" x 36" (61 cm x 91 cm) Drape: 24" x 36" (61 cm x 91 cm) with 4" (10 cm) fenestration Forceps Patient ID Card

- 1: Patient ID Card
 1: Chart Sticker
 1: Patient Information Booklet
 1: Dressing: OpSite® 14 cm x 10 cm
 2: Paper Tape Measure
 1: Tourniquet
 1: Tourniquet
 5: Gauze Pad: 2" x 2" (5 cm x 5 cm)
 6: Gauze Pad: 4" x 4" (10 cm x 10 cm)
 1: Dressing: STATLOCK® Catheter Stabilization Device
 2: Surgical Apparel: Mask
 1: Tape: Steri-Strip® 4
 14 registered trademark of CareFusion or one of its subsidia A registered trademark of CareFusion or one of its subsidiaries.

 A registered trademark of T.J. Smith & Nephew, Limited.

 A registered trademark of C. R. Bard, Inc.

 A registered trademark of 3M Company.

Rx only
Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death. www.teleflex.com/IFU
California Prop. 65

^ WARNING: Cancer and Reproductive Harm.
www.P65Warnings.ca.gov
Not made with natural rubber latex.

Store between 15 - 30°C (59 - 86°E) Avoid freezing and excessive heat.

Store between 15 - 30°C (59 - 86°F). Avoid freezing and excessive heat above 40°C (104°F).

above 40°C (104°F).
Fluid path components are non-pyrogenic.
Contraindications: The PICC is contraindicated wherever there is presence of device related infections or presence of thrombosis in the intended insertion vessel or catheter pathway. Clinical assessment of the patient must be completed to ensure no contraindications exist.

6 Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	Pump Flow Rate†† (mL/hr)	
Distal (18 Ga.)	0.4	380	2450	
Proximal (20 Ga.)	0.4	106	1025	








Arrow International LLC Subsidiary of Teleflex Incorporated 3015 Carrington Mill Blvd. Morrisville, NC 27560 USA



LBL056516 R01 (2022-02)







55 cm catheter length

.018 inch dia. spring-wire guide



^{*} Priming volumes are approximate and are done without accessories.
† Flow rate values are approximate and are determined using deionized water at 100 cm head height.
†† Pump flow rates are determined at maximum pump pressure of 10 psig and represent approximate flow capabilities.