



REF CDC-22122-1A

12 Fr. 2 Lumen 16 cm catheter length .035 inch dia. spring-wire guide



Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume Infusions

Contents:

- 1: Two-Lumen Catheter: 12 Fr. (4.06 mm OD) x 16 cm Radiopaque Polyurethane with Blue FlexTip®, ARROWg+ard® Antimicrobial Surface Treatment

1: Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer

1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle

1: Injection Needle: SafetyGlide™ 1 25 Ga. x 1" (2.54 cm)

1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)

1: Pressure Transduction Probe

1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe

1: Injection Needle: SafetyGlide™ 1 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe

1: Syringe: 3 mL Luer-Lock

1: Tissue Dilator: 10 Fr. (3.5 mm) x 10.2 cm

1: Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm

1: 5 mL 1% Lidocaine HCl Solution and Alcohol Prep

1: 3 mL Applicator 2% CHG and 70% IPA ChlorPrep® One-Step Solution with Hi-Lite Orange™ Tint

1: SharpsAway® Disposal Cup

1: SharpsAway® II Locking Disposal Cup
- 1: Maximal Barrier Drape™ with 4" fenestration

1: Needle Holder

1: Safety Scalpel: #11

1: Checklist/CLIP Sheet

1: Sterile Procedure Sign

1: Medication Label: 1% Lidocaine

2: Gauze Pad: 2" x 2" (5 cm x 5 cm)

5: Gauze Pad: 4" x 4" (10 cm x 10 cm)

1: Surgical Apparel: Impervious Gown

1: Surgical Apparel: Bouffant Cap

1: Surgical Apparel: Mask with Eye Shield

1: Suture: 3-0 Braided Silk with Straight Needle

1: Suture: 3-0 Silk with Curved Needle

1: HemoHopper® Fluid Receptacle
- ¹A trademark of Becton, Dickinson and Company.

²A registered trademark of CareFusion or one of its subsidiaries.

Not made with natural rubber latex.

Store between 20 – 25°C (68 – 77°F).

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.

Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term (≥ 30 days) hemodialysis or for use in patients with thrombosed vessels.


The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/or sulfa drugs.



Priming Volume*

Distal	1.40 mL
Proximal	1.30 mL

*Priming volumes are approximate and are done without accessories.



† In vitro tests performed using 3:1 glycerine/saline at 37°C.

