



## **REF AK-25122-F**

12<sup>Fr.</sup> 2<sup>Lumen</sup>



## Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume Infusions

Contents:

- Two-Lumen Catheter: 12 Fr. (4.06 mm OD) x 20 cm Radiopaque Polyurethane with Blue FlexTip®, 1:
- Arroug+ard® Antimicrobial Surface Treatment, Extension Line Clamps Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 26-3/4" (68 cm) (Straight Soft Tip on One End "J" Tip on 1:
- Other) with Arrow Advancer Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle Injection Needle: 25 Ga. x 1" (2.54 cm) Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) 1:
- 1:
- Pressure Transduction Probe
- Introducer Needle: 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction 1: Syringe
- 1: Injection Needle: 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe Syringe: 3 mL Luer-Lock

- Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm 5 mL 1% Lidocaine HCI Solution and Alcohol Prep 3 mL Applicator 2% CHG and 70% IPA ChloraPrep<sup>⊗1</sup> One-Step Solution with Hi-Lite Orange™ Tint
- 2:
- Dust Cap: Non-Vented SharpsAway® Disposal Cup Drape: 24" x 36" (61 cm x 91 cm) with 4" (10 cm) fenestration
- 1:
- 2 5:
- Gauze Pad: 2" x 2" (5 cm x 5 cm) Gauze Pad: 2" x 2" (5 cm x 5 cm) Suture: 3-0 Braided Silk with Straight Needle 1:
- <sup>1</sup>A registered trademark of CareFusion or one of its subsidiaries.

Rx only

- Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death.

www.P65Warnings.ca.gov

Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F)

- Sterilized using ethylene oxide unless otherwise indicated in the contents list.
- Fluid path components are non-pyrogenic.

Contraindications: The Arrow large-bore two-lumen catheter is not

designed for long-term ( $\geq$  30 days) hemodialysis or for use in patients with thrombosed vessels.

The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/or sulfa drugs.

Priming Volume* (mL)	Flow Rate (mL/min)	Venous Pressure Setting†† (mmHg)
1.6	200	100
Proximal (12 Ga.) 1.5		200
1.5		100
	Volume* (mL)	Volume* Rate (mL/min)   1.6 200 340

\* Priming volumes are approximate and are done without accessories. †† Blood flow rates are approximate and were determined by varying the venous pressure setting from 100 to 200 mmHg in an in vitro experiment using blood analog solution.

## 

LOT

 $\sum$ 

. .







LBL057085 R00 (2022-02)



**REF AK-25122-F** 



.....



**Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume Infusions** 

