





16 cm catheter length .035 inch dia. spring-wire quide

Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume **Infusions**

Contents:

- Two-Lumen Catheter: 12 Fr. (4.06 mm OD) x 16 cm Radiopaque Polyurethane with Blue FlexTip®,
- Arrowg+ard® Antimicrobial Surface Treatment, Extension Line Clamps
 Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End "J" Tip on
- Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle Injection Needle: 25 Ga. x 1" (2.54 cm)
 Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)

- Introducer Needle: 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction
- Injection Needle: 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- Syringe: 3 mL Luer-Lock
- Sylings. Sin Let-Testa.

 Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm

 5 mL 1% Lidocaine HCl Solution and Alcohol Prep

 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®¹ One-Step Solution with Hi-Lite Orange™ Tint
- Dust Cap: Non-Vented SharpsAway® Disposal Cup Drape: 24" x 36" (61 cm x 91 cm) with 4" (10 cm) fenestration

- Scalpel: #11
 Gauze Pad: 2" x 2" (5 cm x 5 cm)
 Gauze Pad: 4" x 4" (10 cm x 10 cm)
 Suture: 3-0 Braided Silk with Straight Needle

¹A registered trademark of CareFusion or one of its subsidiaries.

Lumen	Priming Volume* (mL)	Flow Rate (mL/min)	Venous Pressure Setting†† (mmHg)
Distal (12 Ga.)	1.4	250	100
Proximal (12 Ga.)	1.3	370 250	200 100

* Priming volumes are approximate and are done without accessories. †† Blood flow rates are approximate and were determined by varying the venous pressure setting from 100 to 200 mmHg in an in vitro experiment using blood analog solution.

Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death. California Prop. 65

MARNING: Cancer and Reproductive Harm

www.P65Warnings.ca.gov Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F)

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

Fluid path components are non-pyrogenic.

Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term (≥ 30 days) hemodialysis or for use in patients with thrombosed vessels.

The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/or sulfa drugs.







Arrow International LLC Subsidiary of Teleflex Incorporated 3015 Carrington Mill Blvd. Morrisville, NC 27560 USA



LBL057096 R00 (2022-02)









