





(IPNXXXXXX)



## Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume **Infusions**



- Two-Lumen Catheter: 14 Fr. (4.72 mm OD) x 15 cm Radiopaque Polyurethane with Blue FlexTip®, Arrowg+ard® Antimicrobial Surface Treatment, Extension
- Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow® GlideWheel™ Wire Advance
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer
- Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: Eclipse™1 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
- Injection Needle: Eclipse™1 25 Ga. x 1" (2.54 cm) and 3 mL Luer-Lock Syringe
- Syringe: 10 mL Luer-Lock
- Tissue Dilator: 14 Fr. (4.7 mm) x 14 cm
- Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Maximal Barrier Drape™ with 4" fenestration

- Needle Holder
- Forceps
- Safety Scalpel: #11
- Checklist/CLIP Sheet
- Sterile Procedure Sign
- Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm) Surgical Apparel: Impervious Gown
- Dressing: Tegaderm®2 CHG 3-1/2" x 4-1/2" (8.5 cm x 11.5 cm)
- Transducer Cover: 14 cm x 147 cm
- Surgical Apparel: Mask with Eye Shield
- Surgical Apparel: Bouffant Cap
- Suture: 3-0 Silk with Curved Needle
- HemoHopper® Fluid Receptacle
- <sup>1</sup>A trademark of Becton, Dickinson and Company
- <sup>2</sup>A registered trademark of 3M Company.


## Not made with natural rubber latex.

Store below 25°C (77°F). Avoid excessive heat above 40°C (104°F).

Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term (≥ 30 days) hemodialysis or for use in patients with thrombosed vessels.


The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/ or sulfa drugs.





1.40 mL

Distal



† In vitro tests performed using 3:1 glycerine/saline at 37°C.



LBL071527 R01 (2024-07)